


▲ Figure 17.15 The structure of transfer RNA (tRNA).

Anticodons are conventionally written 3' → 5' to align properly with codons written 5' → 3' (see Figure 17.14). For base pairing, RNA strands must be antiparallel, like DNA. For example, anticodon 3'-AAG-5' pairs with mRNA codon 5'-UUC-3'.

and form a molecule with a three-dimensional structure. Flattened into one plane to clarify this base pairing, a tRNA molecule looks like a cloverleaf (Figure 17.15a). The tRNA actually twists and folds into a compact three-dimensional structure that is roughly L-shaped (Figure 17.15b). The loop extending from one end of the L includes the anticodon, the particular nucleotide triplet that base-pairs to a specific mRNA codon.



▲ Figure 17.16 An aminoacyl-tRNA synthetase joining a specific amino acid to a tRNA. Linkage of the tRNA and amino acid is an endergonic process that occurs at the expense of ATP. The ATP loses two phosphate groups, becoming AMP (adenosine monophosphate).

From the other end of the L-shaped tRNA molecule protrudes its 3' end, which is the attachment site for an amino acid. Thus, the structure of a tRNA molecule fits its function.

The accurate translation of a genetic message requires two instances of molecular recognition. First, a tRNA that binds to an mRNA codon specifying a particular amino acid must carry that amino acid, and no other, to the ribosome. The correct matching up of tRNA and amino acid is carried out by a family of related enzymes called **aminoacyl-tRNA synthetases** (Figure 17.16). The active site of each type of aminoacyl-tRNA


synthetase fits only a specific combination of amino acid and tRNA. (Regions of both the amino acid attachment end and the anticodon end of the tRNA are instrumental in ensuring the specific fit.) There are 20 different synthetases, one for each amino acid; each synthetase is able to bind all the different tRNAs that code for its particular amino acid. The synthetase catalyzes the covalent attachment of the amino acid to its tRNA in a process driven by the hydrolysis of ATP. The resulting aminoacyl tRNA, also called a charged tRNA, is released from the enzyme and is then available to deliver its amino acid to a growing polypeptide chain on a ribosome.

The second instance of molecular recognition is the pairing of the tRNA anticodon with the appropriate mRNA codon. If one tRNA variety existed for each mRNA codon specifying an amino acid, there would be 61 tRNAs (see Figure 17.5). In fact, there are only about 45, signifying that some tRNAs must be able to bind to more than one codon. Such versatility is possible because the rules for base pairing between the third nucleotide base of a codon and the corresponding base of a tRNA anticodon are relaxed compared to those at other codon positions. For example, the nucleotide base U at the 5' end of a tRNA anticodon can pair with either A or G in the third position (at the 3' end) of an mRNA codon. The flexible base pairing at this codon position is called **wobble**. Wobble explains why the synonymous codons for a given amino acid most often differ in their third nucleotide base, but not in the other bases. For example, a tRNA with the anticodon 3'-UCU-5' can base-pair with either the mRNA codon 5'-AGA-3' or 5'-AGG-3', both of which code for arginine (see Figure 17.5).

Ribosomes

Ribosomes facilitate the specific coupling of tRNA anticodons with mRNA codons during protein synthesis. A ribosome consists of a large subunit and a small subunit, each made up of proteins and one or more **ribosomal RNAs (rRNAs)** (Figure 17.17). In eukaryotes, the subunits are made in the nucleolus. Ribosomal RNA genes are transcribed, and the RNA is processed and assembled with proteins imported from the cytoplasm. The resulting ribosomal subunits are then exported via nuclear pores to the cytoplasm. In both bacteria and eukaryotes, large and small subunits join to form a functional ribosome only when they attach to an mRNA molecule. About one-third of the mass of a ribosome is made up of proteins; the rest consists of rRNAs, either three molecules (in bacteria) or four (in eukaryotes). Because most cells contain thousands of ribosomes, rRNA is the most abundant type of cellular RNA.

Although the ribosomes of bacteria and eukaryotes are very similar in structure and function, those of eukaryotes are slightly larger and differ somewhat from bacterial ribosomes in their molecular composition. The differences are medically significant. Certain antibiotic drugs can inactivate bacterial ribosomes without inhibiting the ability of eukaryotic ribosomes to make proteins. These drugs, including tetracycline and streptomycin, are used to combat bacterial infections.



▲ **Figure 17.17** The anatomy of a functioning ribosome.

The structure of a ribosome reflects its function of bringing mRNA together with tRNAs carrying amino acids. In addition to a binding site for mRNA, each ribosome has three binding sites for tRNA, as described in Figure 17.17. The **P site** (**p**eptidyl-tRNA binding site) holds the tRNA carrying the growing polypeptide chain, while the **A site** (**a**minoacyl-tRNA binding site) holds the tRNA carrying the next amino acid to be added to the chain. Discharged tRNAs leave the ribosome from the **E site** (**e**xit site). The ribosome holds the tRNA and mRNA in close proximity and positions the new amino acid for addition to the carboxyl end of the growing polypeptide. It then catalyzes the formation of the peptide bond. As the polypeptide becomes longer, it passes through an *exit tunnel* in the ribosome's large subunit. When the polypeptide is complete, it is released through the exit tunnel.

A lot of evidence strongly supports the hypothesis that rRNA, not protein, is primarily responsible for both the structure and the function of the ribosome. The proteins, which are largely on the exterior, support the shape changes of the rRNA molecules as they carry out catalysis during translation. Ribosomal RNA is the main constituent of the interface between the two subunits and of the A and P sites, and it is the catalyst of peptide bond formation. Thus, a ribosome can be regarded as one colossal ribozyme!

Building a Polypeptide

We can divide translation, the synthesis of a polypeptide chain, into three stages (analogous to those of transcription): initiation, elongation, and termination. All three stages require protein “factors” that aid in the translation process. For certain aspects of chain initiation and elongation, energy is also required. It is provided by the hydrolysis of guanosine triphosphate (GTP), a molecule closely related to ATP.

Ribosome Association and Initiation of Translation


The initiation stage of translation brings together mRNA, a tRNA bearing the first amino acid of the polypeptide, and the two subunits of a ribosome (Figure 17.18). First, a small ribosomal subunit binds to both mRNA and a specific initiator tRNA, which carries the amino acid methionine. In bacteria, the small subunit can bind these two in either order; it binds the mRNA at a specific RNA sequence, just upstream of the start codon, AUG. (Joan Steitz, our Unit Three interviewee, discovered the binding site on the mRNA and showed that complementary base pairing between this site and a ribosomal RNA was involved.) In

eukaryotes, the small subunit, with the initiator tRNA already bound, binds to the 5' cap of the mRNA and then moves, or *scans*, downstream along the mRNA until it reaches the start codon; the initiator tRNA then hydrogen-bonds to the AUG start codon. In either case, the start codon signals the start of translation; this is important because it establishes the codon reading frame for the mRNA.


The union of mRNA, initiator tRNA, and a small ribosomal subunit is followed by the attachment of a large ribosomal subunit, completing the *translation initiation complex*. Proteins called *initiation factors* are required to bring all these components together. The cell also expends energy obtained by hydrolysis of a GTP molecule to form the initiation complex. At the completion of the initiation process, the initiator tRNA sits in the P site of the ribosome, and the vacant A site is ready for the next aminoacyl tRNA. Note that a polypeptide is always synthesized in one direction, from the initial methionine at the amino end, also called the N-terminus, toward the final amino acid at the carboxyl end, also called the C-terminus (see Figure 5.17).

Elongation of the Polypeptide Chain

In the elongation stage of translation, amino acids are added one by one to the previous amino acid at the C-terminus of the growing chain. Each addition involves the participation of several proteins called *elongation factors* and occurs in a three-step cycle described in Figure 17.19. Energy expenditure occurs in the first and third steps. Codon recognition requires hydrolysis of one molecule of GTP, which increases the accuracy and efficiency of this step. One more GTP is hydrolyzed to provide energy for the translocation step.



▲ **Figure 17.18** The initiation of translation.



▲ **Figure 17.19 The elongation cycle of translation.** The hydrolysis of GTP plays an important role in the elongation process. Not shown are the proteins called elongation factors.

The mRNA is moved through the ribosome in one direction only, 5' end first; this is equivalent to the ribosome moving 5' → 3' on the mRNA. The important point is that the ribosome and the mRNA move relative to each other, unidirectionally, codon by codon. The elongation cycle takes less than a tenth of a second in bacteria and is repeated as each amino acid is added to the chain until the polypeptide is completed.


Termination of Translation

The final stage of translation is termination (**Figure 17.20**, on the next page). Elongation continues until a stop codon in the mRNA reaches the A site of the ribosome. The nucleotide base triplets UAG, UAA, and UGA do not code for amino acids but instead act as signals to stop translation. A *release factor*, a protein shaped like an aminoacyl tRNA, binds directly to the stop codon in the A site. The release factor causes the addition of a

water molecule instead of an amino acid to the polypeptide chain. (There are plenty of water molecules available in the aqueous cellular environment.) This reaction breaks (hydrolyzes) the bond between the completed polypeptide and the tRNA in the P site, releasing the polypeptide through the exit tunnel of the ribosome's large subunit. The remainder of the translation assembly then comes apart in a multistep process, aided by other protein factors. Breakdown of the translation assembly requires the hydrolysis of two more GTP molecules.

Polyribosomes

A single ribosome can make an average-sized polypeptide in less than a minute. Typically, however, multiple ribosomes translate an mRNA at the same time; that is, a single mRNA is used to make many copies of a polypeptide simultaneously. Once a ribosome is far enough past the start codon, a second ribosome can attach to the mRNA, eventually resulting in a number



▲ **Figure 17.20 The termination of translation.** Like elongation, termination requires GTP hydrolysis as well as additional protein factors, which are not shown here.

of ribosomes trailing along the mRNA. Such strings of ribosomes, called **polyribosomes** (or *polysomes*), can be seen with an electron microscope (**Figure 17.21**). Polyribosomes are found in both bacterial and eukaryotic cells. They enable a cell to make many copies of a polypeptide very quickly.


Completing and Targeting the Functional Protein

The process of translation is often not sufficient to make a functional protein. In this section, you will learn about modifications that polypeptide chains undergo after the translation process as well as some of the mechanisms used to target completed proteins to specific sites in the cell.


Protein Folding and Post-Translational Modifications

During its synthesis, a polypeptide chain begins to coil and fold spontaneously as a consequence of its amino acid sequence (primary structure), forming a protein with a specific shape: a three-dimensional molecule with secondary and tertiary structure (see Figure 5.20). Thus, a gene determines primary structure, and primary structure in turn determines shape. In many cases, a chaperone protein (chaperonin) helps the polypeptide fold correctly (see Figure 5.23).

Additional steps—*post-translational modifications*—may be required before the protein can begin doing its particular job in the cell. Certain amino acids may be chemically modified by the attachment of sugars, lipids, phosphate groups, or other additions. Enzymes may remove one or more amino acids from the leading (amino) end of the polypeptide chain. In some cases, a polypeptide chain may be enzymatically cleaved into two or more pieces. For example, the protein insulin is first synthesized as a single polypeptide chain but becomes active only after an enzyme cuts out a central part of the chain,



(a) An mRNA molecule is generally translated simultaneously by several ribosomes in clusters called polyribosomes.



(b) This micrograph shows a large polyribosome in a bacterial cell. Growing polypeptides are not visible here (TEM).

▲ **Figure 17.21 Polyribosomes.**

leaving a protein made up of two polypeptide chains connected by disulfide bridges. In other cases, two or more polypeptides that are synthesized separately may come together, becoming the subunits of a protein that has quaternary structure. A familiar example is hemoglobin (see Figure 5.20).